

Electronic Smoking Devices and Secondhand Aerosol

Electronic smoking devices (or ESDs), which are often called **e-cigarettes**, heat and vaporize a solution that typically contains nicotine. The devices are metal or plastic tubes that contain a cartridge filled with a liquid that is vaporized by a battery-powered heating element. The aerosol is inhaled by the user when they draw on the device, as they would a regular tobacco cigarette, and the user exhales the aerosol into the environment.

“If you are around somebody who is using e-cigarettes, you are breathing an aerosol of exhaled nicotine, ultra-fine particles, volatile organic compounds, and other toxins.” Dr. Stanton Glantz, Director for the Center for Tobacco Control Research and Education at the University of California, San Francisco.

Current Legislative Landscape

- As of January 1, 2024, [1,054 municipalities, plus 27 states, commonwealths, and territories](#) restrict e-cigarette use in 100% smokefree venues.

Constituents of Secondhand Aerosol

Electronic smoking devices (ESDs) do not just emit “harmless water vapor.” **Secondhand aerosol (incorrectly called vapor by the industry) from ESDs contains nicotine, ultrafine particles and low levels of toxins** that are known to cause cancer.

- ESD aerosol is made up of a high concentration of ultrafine particles, and the particle concentration is higher than in conventional tobacco cigarette smoke.ⁱ
- Exposure to fine and ultrafine particles may exacerbate respiratory ailments like asthma, and constrict arteries which could trigger a heart attack.ⁱⁱ
- ESD aerosol particles are smaller than 1000 nanometers, which is a similar size to tobacco smoke and diesel engine smoke, and bystanders can be exposed to this aerosol. “The exact size distribution depends on the chemical composition of the electronic cigarette liquid, the e-cigarette device operation, and user vaping preferences.”ⁱⁱⁱ
- At least 10 chemicals identified in ESD aerosol are on California’s Proposition 65 list of carcinogens and reproductive toxins, also known as the [Safe Drinking Water and Toxic Enforcement Act of 1986](#). The compounds that have already been identified in ESD aerosol include: **Acetaldehyde (MS), Benzene (SS), Cadmium (MS), Formaldehyde (MS,SS), Isoprene (SS), Lead (MS), Nickel (MS), Nicotine (MS, SS), N-Nitrosornicotine (MS, SS), Toluene (MS, SS).**^{iv,v}
- ESDs contain and emit propylene glycol**, a chemical that is used as a base in ESD solution and is one of the primary components in the aerosol emitted by ESDs.
 - Short term exposure causes eye, throat, and airway irritation.^{vi}
 - Long term inhalation exposure can result in children developing asthma.^{vii}
- Even though propylene glycol is FDA approved for use in some products, the inhalation of vaporized nicotine in propylene glycol is not. Some studies show that heating propylene glycol changes its chemical composition, producing small amounts of propylene oxide, a known carcinogen.^{viii}

- There are **metals in ESD aerosol, including chromium, nickel, and tin nanoparticles**.^{ix}
- FDA scientists found detectable levels of carcinogenic tobacco-specific nitrosamines in ESD aerosol.^x
- People exposed to ESD aerosol absorb nicotine (measured as cotinine), with one study showing levels comparable to passive smokers.^{xi}
- **Diethylene Glycol**, a poisonous organic compound, was also detected in ESD aerosol.^{xii}
- **Exhaled ESD aerosol contained propylene glycol, glycerol, flavorings, and nicotine, along with acetone, formaldehyde, acetaldehyde, propanal, diacetyl, and triacetyl**.^{xiii}
- Many of the elements identified in the aerosol are known to **cause respiratory distress and disease**. The aerosol contained particles >1 µm comprised of tin, silver, iron, nickel, aluminum, and silicate and nanoparticles (<100 nm) of tin, chromium and nickel. The concentrations of nine of eleven elements in ESD aerosol were higher than or equal to the corresponding concentrations in conventional cigarette smoke.^{xiv}
- ESDs cause exposure to different chemicals than found in conventional cigarettes and there is a need for risk evaluation for both primary and passive exposure to the aerosol in smokers and nonsmokers.^{xv}
- Short term use of ESD has been shown to increase respiratory resistance and impair lung function, which may result in difficulty breathing.^{xvi}
- The first study to look at exposure to aerosol from ESDs in real-use conditions found that non-smokers who were exposed to conventional cigarette smoke and ESD aerosol absorbed similar levels of nicotine.^{xvii}
- The American Society of Heating, Refrigerating, and Air Conditioning Engineers (ASHRAE) has concluded that ESDs emit harmful chemicals into the air and need to be regulated in the same manner as tobacco smoking. The “E-cigarettes do not produce a vapor (gas), but rather a dense visible aerosol of liquid sub-micron droplets consisting of glycols, nicotine, and other chemicals, some of which are carcinogenic (e.g., formaldehyde, metals like cadmium, lead, & nickel, and nitrosamines).”^{xviii}
- ESD aerosol is a source of high doses of particles being deposited in the human respiratory system.^{xix}
- ESD exposure damages lung tissues. Human lung cells that are exposed to ESD aerosol and flavorings—especially cinnamon—are show increased oxidative stress and inflammatory responses.^{xx}
- Concentrations of formaldehyde are higher than concentrations of nicotine in some samples of ESD aerosol. Formaldehyde is created when propylene glycol and glycerol are heated to temperatures reached by commercially available ESDs operating at high voltage.^{xxi}
- Flavorings are a largely unrecognized potential hazard of ESDs. Diacetyl and acetyl propionyl are present in many sweet-flavored ESDs, and are approved by the FDA for food use (ingestion), but are not evaluated and approved for heating and inhalation, and are associated with respiratory disease when inhaled.^{xxii} High doses of diacetyl, used to flavor buttered popcorn, have been shown to cause acute-onset bronchiolitis obliterans, a severe and irreversible obstructive lung disease when inhaled by workers exposed to particulate aerosolized flavorings containing diacetyl.^{xxiii} Therefore, these chemicals cannot be deemed “generally recognized as safe” for inhalation.

- Nanoparticles in ESD aerosol are much smaller than the particles in tobacco smoke and are present in much higher concentrations. Toxic chemicals attached to nanoparticles may have greater adverse health effects than when these toxins are attached to larger tobacco smoke particles.^{xxiv} Nanoparticles are more easily and deeply breathed into the lungs of the user and bystander.
- ESD aerosols contain carbonyls at levels which can have cardiovascular toxicity. While ESD aerosol has lower levels of toxins than tobacco smoke, toxins from the aerosol may still have a significant cardiovascular impact because cardiovascular disease has a nonlinear dose-response, which means that high risk is possible with relatively low exposure.^{xxv}
- Human lung cells exposed to ESD aerosol and copper nanoparticles show signs of inflammatory stress and DNA fragmentation.^{xxvi}
- ESD use alters the physical appearance of airways and may impact the development of chronic lung disease. The airways of people who use ESDs appear redder than the airways of both people who smoke and nonsmokers.^{xxvii}
- ESDs that operate using a single-coil heating element produce much higher levels of toxins than double-coil devices across different e-liquids. Double-coil devices produce aerosol at lower temperatures while single-coil devices produce aerosol at higher temperatures.^{xxviii}
- Daily ESD users have double the risk of heart attack, and the dual use of ESDs and conventional cigarettes—which is the most common use pattern among ESD users—is more dangerous than using either product alone.^{xxix}
- There is a risk of thirdhand exposure to nicotine released from ESD aerosol that deposits on indoor surfaces.^{xxx}
- Chemicals from ESDs can drift through multi-unit buildings and deposit on surfaces in spaces where ESDs are not being used.^{xxxi} Overall, ESDs are a new source of **Volatile Organic Compounds (VOCs) and ultrafine/fine particles in the indoor environment**, thus resulting in “passive vaping.”^{xxxii}
- The World Health Organization (WHO) recommends that ESDs not be used indoors, especially in smokefree environments, in order to minimize the risk to bystanders of breathing in the aerosol emitted by the devices and to avoid undermining the enforcement of smokefree laws.^{xxxiii}
- The National Institute for Occupational Safety and Health (NIOSH) recommends that employers “establish and maintain smoke-free workplaces that protect those in workplaces from involuntary, secondhand exposures to tobacco smoke and airborne emissions from e-cigarettes and other electronic nicotine delivery systems.”^{xxxiv}
- The American Industrial Hygiene Association (AIHA) also recommends that ESDs be included in smokefree laws: **“Because e-cigarettes are a potential source of pollutants (such as airborne nicotine, flavorings, and thermal degradation products), their use in the indoor environment should be restricted, consistent with current smoking bans, until and unless research documents that they will not significantly increase the risk of adverse health effects to room occupants.”**^{xxxv}
- The American Public Health Association adopted a resolution, “Supporting Regulation of Electronic Cigarettes,” that outlines seven action steps including, “States and municipalities [should] enact and enforce laws...prohibiting the use of e-cigarettes in all enclosed areas of public access and places of employment. These standards should be incorporated into existing clean indoor air laws.”^{xxxvi}

- The American Association for Cancer Research and the American Society of Clinical Oncology supports prohibiting the use of ESDs in smokefree spaces until the safety of second- and thirdhand aerosol exposure is established.^{xxxvii}

ESD aerosol is a new source of pollution and toxins being emitted into the environment. We do not know the long-term health effects of ESD use and although the industry marketing of the product implies that these products are harmless, the aerosol that ESD emit is not purely water vapor.

May be reprinted with appropriate attribution to the American Nonsmokers' Rights Foundation, © 2024

For more information, contact us at info@no-smoke.org or call 510-841-3032

REFERENCES

-
- ⁱ Fuoco, F.C.; Buonanno, G.; Stabile, L.; Vigo, P., "[Influential parameters on particle concentration and size distribution in the mainstream of e-cigarettes](#)," *Environmental Pollution* 184: 523-529, January 2014.
- ⁱⁱ Grana, R.; Benowitz, N.; Glantz, S., "[Background Paper on E-cigarettes](#)," Center for Tobacco Control Research and Education, University of California, San Francisco and WHO Collaborating Center on Tobacco Control. December 2013.
- ⁱⁱⁱ Thornburg, J.; Malloy, Q.; Cho, S.; Studabaker, W.; Lee, Y.O., "[Exhaled electronic cigarette emissions: what's your secondhand exposure?](#)," *RTI Press Research Brief*: 1-4, March 2015.
- ^{iv} Goniewicz, M.L.; Knysak, J.; Gawron, M.; Kosmider, L.; Sobczak, A.; Kurek, J.; Prokopowicz, A.; Jablonska-Czapla, M.; Rosik-Dulewska, C.; Havel, C.; Jacob, P.; Benowitz, N., "[Levels of selected carcinogens and toxicants in vapour from electronic cigarettes](#)," *Tobacco Control* [Online ahead of print], March 6, 2013.
- ^v Williams, M.; Villarreal, A.; Bozhilov, K.; Lin, S.; Talbot, P., "[Metal and silicate particles including nanoparticles are present in electronic cigarette cartomizer fluid and aerosol](#)," *PLoS ONE* 8(3): e57987, March 20, 2013.
- ^{vi} Wieslander, G.; Norbäck, D.; Lindgren, T., "[Experimental exposure to propylene glycol mist in aviation emergency training: acute ocular and respiratory effects](#)," *Occupational and Environmental Medicine* 58:10 649-655, 2001.
- ^{vii} Choi, H.; Schmidbauer, N.; Spengler, J.; Bornehag, C., "[Sources of Propylene Glycol and Glycol Ethers in Air at Home](#)," *International Journal of Environmental Research and Public Health* 7(12): 4213–4237, December 2010.
- ^{viii} Henderson, TR; Clark, CR; Marshall, TC; Hanson, RL; & Hobbs, CH. "[Heat degradation studies of solar heat transfer fluids](#)," *Solar Energy*, 27, 121-128. 1981.
- ^{ix} Williams, M.; Villarreal, A.; Bozhilov, K.; Lin, S.; Talbot, P., "[Metal and silicate particles including nanoparticles are present in electronic cigarette cartomizer fluid and aerosol](#)," *PLoS ONE* 8(3): e57987, March 20, 2013.
- ^x Westenberger, B.J., "Evaluation of e-cigarettes," St. Louis, MO: U.S. Department of Health and Human Services (DHHS), Food and Drug Administration (FDA), Center for Drug Evaluation and Research, Division of Pharmaceutical Analysis, May 4, 2009.
- ^{xi} Flouris, A.D.; Chorti, M.S.; Pouliant, K.P.; Jamurtas, A.Z.; Kostikas, K.; Tzatzarakis, M.N.; Wallace, H.A.; Tsatsaki, A.M.; Koutedakis, Y., "[Acute impact of active and passive electronic cigarette smoking on serum cotinine and lung function](#)," *Inhalation Toxicology* 25(2): 91-101, February 2013.
- ^{xii} Westenberger, B.J., "Evaluation of e-cigarettes," St. Louis, MO: U.S. Department of Health and Human Services (DHHS), Food and Drug Administration (FDA), Center for Drug Evaluation and Research, Division of Pharmaceutical Analysis, May 4, 2009.
- ^{xiii} Schripp, T.; Markewitz, D.; Uhde, E.; Salthammer, T., "[Does e-cigarette consumption cause passive vaping?](#)," *Indoor Air* 23(1): 25-31, February 2013.
- ^{xiv} Williams, M.; Villarreal, A.; Bozhilov, K.; Lin, S.; Talbot, P., "[Metal and silicate particles including nanoparticles are present in electronic cigarette cartomizer fluid and aerosol](#)," *PLoS ONE* 8(3): e57987, March 20, 2013.
- ^{xv} Pellegrino, R.M.; Tighino, B.; Mangiaracina, G.; Marani, A.; Vitali, M.; Protano, C.; Osborn, J.F.; Cattaruzza, M.S., "[Electronic cigarettes: an evaluation of exposure to chemicals and fine particulate matter \(PM\)](#)," *Annali Di Igiene* 24(4):279-88, July-August 2012.
- ^{xvi} Vardavas, C.I.; Anagnostopoulos, N.; Kougias, M.; Evangelopoulou, V.; Connolly, G.N.; Behrakis, P.K., "[Short-term pulmonary effects of using an electronic cigarette: impact on respiratory flow resistance, impedance, and exhaled nitric oxide](#)," *Chest* 141(6): 1400-1406, June 2012..

- xvii ["Cigarettes vs. e-cigarettes: Passive exposure at home measured by means of airborne marker and biomarkers."](#) *Environmental Research*, Volume 135, November 2014.
- xviii Offermann, Bud. ["The Hazards of E-Cigarettes."](#) ASHRAE Journal, June 2014.
- xix Manigrasso, M.; Buonanno, G.; Fuoco, F.C.; Stabile, L.; Avino, P., ["Aerosol deposition doses in the human respiratory tree of electronic cigarette smokers,"](#) *Environmental Pollution* 196: 257-267, January 2015.
- xx Lerner CA, Sundar IK, Yao H, Gerloff J, Ossip DJ, McIntosh S, et al. ["Vapors Produced by Electronic Cigarettes and E-Juices with Flavorings Induce Toxicity, Oxidative Stress, and Inflammatory Response in Lung Epithelial Cells and in Mouse Lung,"](#) *PLoS ONE* 10(2): e0116732, February 6, 2015.
- xxi Jensen, R.P.; Luo, W.; Pankow, J.F.; Strongin, R.M.; Peyton, D.H., ["Hidden formaldehyde in e-cigarette aerosols,"](#) *New England Journal of Medicine* 372: 392-394, January 22, 2015.
- xxii Konstantinos E. Farsalinos, KE; Kistler, KA; Gilman, G; Voudris, V., ["Evaluation of electronic cigarette liquids and aerosol for the presence of selected inhalation toxins,"](#) *Nicotine and Tobacco Research* 17(2): 168-174, February 2015.
- xxiii ["Preventing Lung Disease in Workers: Who Use or Make Flavorings,"](#) National Institute for Occupational Safety and Health. DHHS (NIOSH) Publication Number 2004–110, December 2003.
- xxiv Mikheev, V.B.; Brinkman, M.C.; Granville, C.A.; Gordon, S.M.; Clark, P.I., ["Real-time measurement of electronic cigarette aerosol size distribution and metals content analysis,"](#) *Nicotine and Tobacco Research* [Online ahead of print], May 4, 2016.
- xxv Bhatnagar, A. ["E-cigarettes and Cardiovascular Disease Risk: Evaluation of Evidence, Policy Implications, and Recommendations,"](#) *Current Cardiovascular Risk Report* 10: 24, July 2016.
- xxvi Lerner, C.A.; Rutagarama, P.; Ahmad, T.; Sundar, I.K.; Elder, A.; Rahman, I., ["Electronic cigarette aerosols and copper nanoparticles induce mitochondrial stress and promote DNA fragmentation in lung fibroblasts,"](#) *Biochemical and Biophysical Research Communications* 477(4): 620-625, September 2, 2016.
- xxvii Ghosh, A., Coakley, R. C., Mascenik, T., Rowell, T. R., Davis, E. S., Rogers, K., ... Tarran, R. (2018). Chronic e-cigarette exposure alters the human bronchial epithelial proteome. *American Journal of Respiratory and Critical Care Medicine*, 198(1), 67-76. DOI: 10.1164/rccm.201710-2033OC
- xxviii Sleiman, et al. ["Emissions from Electronic Cigarettes: Key Parameters Affecting the Release of Harmful Chemicals."](#) Lawrence Berkeley National Laboratory. Journal of Environmental Science Technology. July 27, 2016.
- xxix Talal Alzahrani, Ivan Pena, Nardos Temesgen, Stanton A. Glantz. Association Between Electronic Cigarette Use and Myocardial Infarction. *Am J Prev Med* 2018; Published online ahead of print 22-AUG-2018 DOI information: 10.1016/j.amepre.2018.05.004.
- xxx Goniewicz, M.L.; Lee, L., ["Electronic cigarettes are a source of thirdhand exposure to nicotine,"](#) *Nicotine and Tobacco Research* [Online ahead of print], August 30, 2014.
- xxxi Khachatoorian C, Jacob III P, Benowitz NL, et al. ["Electronic cigarette chemicals transfer from a vape shop to a nearby business in a multiple-tenant retail building."](#) Tobacco Control Published Online First: 29 August 2018. doi: 10.1136/tobaccocontrol-2018-054316.
- xxxii Schripp, T.; Markewitz, D.; Uhde, E.; Salthammer, T., ["Does e-cigarette consumption cause passive vaping?,"](#) *Indoor Air* 23(1): 25-31, February 2013.
- xxxiii World Health Organization (WHO), ["Electronic nicotine delivery systems,"](#) World Health Organization (WHO), 2014.
- xxxiv ["Promoting Health and Preventing Disease and Injury Through Workplace Tobacco Policies,"](#) National Institute for Occupational Safety and Health (NIOSH), Current Intelligence Bulletin 67, April 2015.
- xxxv ["White Paper: Electronic Cigarettes in the Indoor Environment,"](#) American Industrial Hygiene Association, October 19, 2014.
- xxxvi American Public Health Association. ["Supporting regulation of electronic cigarettes,"](#) APHA Policy Statement Number 20149, November 18, 2014.
- xxxvii Brandon, T.H.; Goniewicz, M.L.; Hanna, N.H.; Hatsukami, D.K.; Herbst, R.S.; Hobin, J.A.; Ostroff, J.S.; Shields, P.G.; Toll, B.A.; Tyne, C.A.; Viswanath, K.; Warren, G.W., ["Electronic nicotine delivery systems: a policy statement from the American Association for Cancer Research and the American Society of Clinical Oncology,"](#) *Journal of Clinical Oncology* [Online ahead of print], January 8, 2015.